INTRAMED Plus
infusion services

RITUXAN®
For GPA or MPA

INFUSION & MEDICAL CENTER

1. Patient Name ______________________ **DOB** __________ **Patient Phone/Cell #** ______________

Patient demographic and insurance information to be faxed with Infusion Order Form

2. Medical Information (Please select primary diagnosis and complete ICD10 Code):

Primary Diagnosis: ______Granulomatosis with Polyangiitis (GPA) ICD-10 Code: M31.30 _____

______Microscopic Polyangiitis (MPA) ICD-10 Code: M31.7 ______

______Other: ICD-10 Code: __________

Allergies:____(or attach list)

3. Clinical Information – Please fax with Infusion Order Form:

Patient
Weight: __________ lbs.
Height:___________ in.

- Clinical MD Notes, labs, test supporting primary diagnosis
- Pre-Screening Documentation
  Hepatitis B Screening Results (including Hep B surface antigen & Total Hep B Core Antibody)
- Previous Drug Therapy History, including therapies trialed and or failed and date of last infusion:
  - ❑ Previous biologic therapies: ________________________________Date:______________
  - ❑ Washout period of ________ weeks desired prior to the initiation of this ordered therapy
- Infusion Center – Lab Orders (Check for Infusion Center to Manage):
  - ❑ Obtain CBC with diff and platelets every ___________
- Corticosteroid Regimen: Has your patient started on a steroid regimen prior to receiving Rituxan? ❑ Yes ❑ No
  If yes, provide corticosteroid regimen: ______________________________________

RITUXAN® (rituximab) J Code: J9312

4. Drug Order: Administer Rituxan IV as per the below parameters:

Induction Dose: ❑ 375 mg/m² once weekly x 4 weeks or ❑ Other:___________

Maintenance Dose: ____

Pre-Medication Orders: Administer Acetaminophen 650 mg PO; Diphenhydramine 50 mg PO orally 30 minutes prior to infusion and adjust to patient's needs PLUS

❑ Induction Steroid Therapy: Methylprednisolone 1000mg IV Daily x 3 doses prior to Rituxan therapy or adjusted according to prior steroid dosing regimen.

If induction steroid therapy is completed, Methylprednisolone 100 mg IV 30 mins prior to infusion.

Other:____

Adverse Drug Reaction Protocol: Manage any adverse reaction that may occur per approved ADR Protocol.

By signing this form and utilizing these services, I am authorizing Intramed Plus to serve as my prior authorization agent with medical and pharmacy insurance providers.

5. Physician Signature: ________________________ / ________________________ Date: __________

Dispense as written Substitution permitted

Printed Physician's Name:______________________________Contact Phone #: __________________

FAX ALL INFORMATION CENTRAL FAX **803.999.1754**	**INFUSION CENTER LOCATIONS** COLUMBIA CHARLESTON GREENVILLE CENTRAL INTAKE PHONE 803.999.1760